

Toxicology Excellence For Risk Assessment 1250 Ohio Pike, Suite #197 Cincinnati, Ohio 45102-1239

June 3, 2019

To: Science Advisory Board, U.S. Environmental Protection Agency

From: Michael L. Dourson, PhD., DABT, FATS, FSRA
Director of Science, Toxicology Excellence for Risk Assessment (TERA)
President, Toxicology Education Foundation

Re: Public Comments

Dear Dr. Honeycutt and Colleagues,

I offer the following comments on issues before your esteemed committee.

- Although it no doubt can be improved and abuse must be safeguarded, the **Science Transparency Rule** is necessary from a risk assessment perspective. Why? An example may be instructive. We currently have one chemical in judicial review based in large part on an observational epidemiology study suggesting IQ deficits in children. If these effects were true, this would be of grave concern. However, the study is in contrast to the wealth of other guideline-study data and it's suggested hypothesis has been tested in experimental animals and found not to be supported. Moreover, the published study is missing up to 35% of the data stated to be otherwise available as demonstrated in a recent SOT poster (attached) and the authors have not released their data, even redacted, to EPA, despite being asked twice, despite the study being publicly funded in part, and despite EPA's ability to handle otherwise confidential data routinely. Determining a risk assessment position with such data is highly uncertain and easily challenged by other scientists. Not using data in such situations is a reasonable policy decision.
- EPA has published progressive Guidelines for Carcinogen (2005) and Noncancer (2002)
 Assessment and is acclaimed by all to be a leader in these areas as demonstrated by extensive use and citation. However, risk assessment science has matured from the date of these last publications. Concepts such as Data-derived extrapolation factors (EPA, 2014), Bayesian Benchmark Dose (BMD), risk above the safe dose, high throughput toxicity screens, adverse outcome pathway (AOPs), and new thinking on mode of action (MOA) all can, and should, be integrated into revised guidelines. Several of these topics can easily and quickly be incorporated. Other topics will likely require additional study, debate, and judgment. See a recent poster at the NAS (2019) for one such consideration on dual MOA/AOP for the cancer findings of acrylamide (attached).
- **Per- and Polyfluoroalkyl Substances (PFAS) Action Plan** is important and as such should incorporate any newly available data. Recent findings of PFOA administration to 43 cancer patients with good renal clearance and outstanding kinetic follow-up appear to contradict observational studies of long half lives in humans and suggest different approaches to the highly disparate (~750-fold) safe dose assessment of different governments. EPA can take the lead here in both a revised safe dose assessment and in research on human PFOA/S kinetic clearance to

- address a more appropriate animal to human extrapolation. A recent paper submitted on this topic can be accessed at: https://www.tera.org/about/news.html (see 5-14-19 story).
- EPA's Proposed Waters of the U.S. Rule is important and more or less consistent with the definition of *navigable waters* in the act. Although the desire to extend the rule to vernal waters, swamps, or marshes as some suggest is not unreasonable from several viewpoints, it would necessitate the application of ambient water quality criteria to such waters. The development of these criteria assumes that humans drink 2 liters of water per day and eat 6 to 20 grams of fish per day from such waters. From a practical occurrence and risk assessment perspective this makes little sense. The SAB might consider this risk perspective in their discussions as appropriate.


If time allows, I would be happy to answer any questions.

Sincerely,

1

Michael L. Dourson, Ph.D., DABT, FATS, FSRA Director of Science

Independent ••• Non-Profit ••• Science A 501c3 Environmental Science NGO





Evidence-Based Dose Response Assessment for Thyroid Tumorigenesis: A tale of two MOAs.


and what can be done about it

Hazard Identification 1

- Actylatnisk met skenet te onner maret in hannan (Ma al., 2007), Newson et al., 2007, Mancel & Adams, 2005).
- Six king-nava experiments in east show turnan (kikinasa et al 1986; Friedman et al., 1995; and NTP, 2012). All experiment used the same strain of tota.
- lable in mace (NTP, 2912). * Two long terms stadius also
- Francisco serio seriodore, are aleso arraitatón in mice send cuto f glycinhamide, a koncern metaboliste est acrytamide, se visro.

Mode of Action 1

- Arcylandia is geostavit, that on decidyly mangania. Arcylandia also execute provide stimulation and collative service, the inter of which can lead to example on and collative service, the inter of which can lead to examinate and celebrary geometric exposure are not show response conventions—and you see a doors injustic exposure are not show response conventions—and you see a doors injustic than the top of the collection of



Dose Response Assessment 1

- EPA (2005) suggests "decouping" data when servent of action occur in definition parts of the deep response. These.
- The control of the
 - Thus, a 2% BMR is correlectably within the interpolation allowing for a stable extension in the point of departmen.

Propose 28, Multiploage model Ribad to provided all thyroid branch date, stronger 1918, the change is dolpe branch date, stronger 1918, the stronger is dolpe branch to but and high done regions (Dourson et al., 2MS). Figure 6at, S S S S

Dose Response Assessment 2

Low dose to vivo genotice low-dose burners

Mode of Action 2

Hazard Identification 2

Acrybamide is carpmently found in various foods.

khing EPA (2007) gaidelines, we compared different math models to til these "decoupled" lists.

Companions with Table 2 of EPA (1908)

Wast eensitive study a only the most patent (bes Appendix to this preventation)

We identify a connece stape factor of 6.000 (magical days). See how doze, and a \$110 of 6.05 to 0.02 magical day by high doze

 We "downight" those MOAs and determine that the probit execute best reflects the underlying "documpled" biology. Book mantapresiety & grewith-ethnetistion occur; metagore determines fore-dose; growth nitroulation at higher desert Canuse studies in pass region increases in many types of tunsors. We find that disposed tunsors are most sensitive because they are represented, receiving may pointed.

of this manipul, adid by being an other concentration to be of sufficient quality (California, 2012). A real association that the A real association of programs of volume and the concentration and the deterministing the suit in human scholars catelion that are suggesteritely. A seministic market such services that are bound in comparable oursiles of the same or other spoored/strain. Princine. A semines that phase effects in one species that are expected to be found in human.

Figure 3s. Protes model filted to ported all frynaid tumor data. showing differing shopes between the low and high dose regions. (Dostreen et A., 2018, Figure 6b).

For Conference of the Conferen

Starch tumora approarance is mare nounsistent with measurem or tumor formataion fasts are not formed after a respective. Those observations also measure it a unified by the direct measurisms canning aid of fits cannow in these experiments.

Although, scienties have not identified say, themisel that has caused thyread assures to huranes and the rat dystud is different form the human thyread, we contable that these not thyreds true not subjectly to the subject of the problem of the problem to the subject of the problem of the problem.

Whe senderbook is more contributionary some...

We analyze from the following:

- new more was analyzed by the following:

- new more was anythmetely to the form made of more and

- new more was anythmetely for the form and the form th

Previous assessabents were fromd to be gene "securing level" assessments.

Background

Kory decementations for evaluating tention responses are:

 consistently elegency that can be entroposity opticises tention to the control of the

Them, both a munipum: formation are likely to a

The type of thyroid tumour formed in sute is generally recognize as resulting from growth strongleton assive routstines, and these roodes of action alon sported in humans.

C 1284

Yumans evoked by actylamide exposure were generally brough, economic law in life, and were mean other in hermanily-actors organe, in all the rat experiments.

Mode of Action 3


Hazard Identification 3

Thyroid summin in rath orquines to acrylamide wore man earlielically significant in all six experiments (Table 1).

Dose Response Assessment 3

Table 2. Key events, related evidence and plausibility of tumor modes of action.

- The problet model Bonchenard, done (BMD) of 2% curre risks is 0.81 mpg legyle, paracited out in a drope of 10.52 mpg legyle, paracited out in a drope of 10.52 mpg legyle, paracited out in a drope of 10.52 mpg legyle, dones in the outer of 10.52 mpg legyle out in the outer of 10.52 mpg legyle of 10.52 mpg legyle, out out of 10.52 mpg legyle of 10.52 mpg legyle, outside of 10.52 mpg legyle of 10.52 mpg legyle outside out of 10.52 mpg legyle outside in the 10.52 mpg legyle outside of 10.52 mpg legyle outside in the 10.52 mpg legyle outside in the 10.52 mpg legyle outside in particular outside in the 10.52 mpg legyle in the 10
 - The adjusted alope factor is $0.050~(\mathrm{mg/kg-day})^{1.5}$



- Mission is to support the perfection of public health by
 - Bovedoping, reviewing and commonituding risk
 associations values and analyses:

 Improveing risk methods throught research; and
 Edwarding risk association; managers, and the publish
- TERA is a 2010 houseoft

Figure 1. Thyroid Sumers Plubed

Party on	RMB-62	がお野様があ	SPARM #5	S a 850% 13
Studyed Sprands	6.83	100	100	6034
Pincked syads	6.43	4,803	85.3	8.634
Postedal	100	\$283	8.3	9.038
	91 094	经有限的证据	W400.13	5F or \$50,000, 1.8
Period forage:	80	0.867	100	8,686
Packs sale	6.7	637.8	w.	9/8/9
Parketsk	F-	6,857	3.1	0.083

***	BARBL	18.88		erkin .		20,93		18.00		2000
	Model	Sylventers		Machine		Profes		Websell.		0.000
	Been, Not	100	Season	Live day	passing appears	100	Desirate Control	*	protect	-
00000	-	-	2000	(ME	500	255	AND S	S012	005	e R
				-	****	200	in the second	***		
id. Lattie 6).		PADA #1 SF or BADEL #1	404	6.974	6.028	200	5F or \$50,000, 14	1.04%	0.079	

of 2% BMSS and BMSS, from various funed fryend furnors in rate & projected stops facture. SE RESIDEA SE RESPONDADO 8363 625 maga 8.9 \$1400 Bully 8,400 2.2 6,925 0,93 6,004 2,8 6-828 6-75 K-62K 6-2

- Company of the Compan